
Liquid-crystalline medium

the inventor is related to a light intight alone weight and the importance of the section of the integral of the section of th



and at least the tip unit into the inflate w

$$R^{2}$$
 H \rightarrow alkenyl

Fig. F. and F. are each, independently it she enother, an alkylor alkenyloradical harring up to take a structure in a modern cell of the control of the cont

in the Millian William William

are not conket threatly to be another.

the dethological and result of the inflame swetching. Bug layer

The principle of electrically through the first framework the Ethorism of alternatively laborated the first time in 1971 (M.F. swillerkel and E. Fahrenschen, "Deformation or nematical treatm of approximation of the first fields" Appl. Phys. Lett. 18 1071 (Birs. 1971) This was it is well symptoms by U.F. Fann Appl. Phys. Lett. 19 22 (Appl. Phys. Lett. 19 22) Appl. Phys. Lett. 19 22 (Appl. Phys. Lett. 19 22) Appl. Phys. Lett. 19 22 (Appl. Phys. Lett. 19 22) Appl. Phys. Lett. 19 22 (Appl. Phys. Ref. Phys. Lett. 19 22) Appl. Phys. Ref. Phys. Lett. 20 22 (Appl. Phys. Ref. Phys. Phys.

The papers by J. Fried and E. Tlene SID Si Disest Terim. Papers I and E. Schail E. Disest Terim. Papers I and E. Schail ESID SI Disest Terim. Papers I and E. Schail ESID SI Disest Terim. Papers I and E. Data may show that liquid stylealline phases must have in measured in the values of rethe ratio between the elastic penatants E. F. high values of rethe prival anisotropy As of from Life to Sensor prival entries to be much able of rehish information singley elements a section the E.B. effect. Flection prival implicy elements based on the E.B. effect. Flection in the prival implicate accommend. Discourtable by a gative I prival organic medical tanks also be used in Hisplays willight medical tanks as the used in Hisplays willight them.

Is affined a long to the control of the control of

in the second of the second of

In the contract the open place of the contract of the contract

Matrix (s.p. i reported limpleyer are an who Hom line of elements which has be used to individual switching of the individual pixels are, for example, active elements line, transistors. This is then referred to as an magnite matrix, and a distinction was be made between two types:

- 1. M.J. meral kije semiseniko i trangibi is en a ... silisen water ab kubarrate.
 - 11. Inches in the state of the state of the state of the state.
- In the case of type 1, the electron prical effect used is usually signatic scarrening a the mesh hist effect. The use of schole hypital price is as the significant material limits the diagraph size, while even in that assembly of case use part in plays results in problems.

ling the company of the respondence of the company of the company

A file general designation formation for the state of the

inventure EqR or is to receive the well-wase on the latter terminally.

Include that is the angles while the unities of he distributed plate there is the transparent your erelected described the THI universe with the wine of the pixel electrodes the THI universe whall and has wire also not appeared effects in the instead of the pixel electrodes. The instead of the instead of the pixel electrodes are distributed to fully the companies of the displayer in which a matrix of red, are encounting the electrodes and a section of the expense of the electrodes are displayed in which a way that electrodes electron is a least of protection with the electrodes.

1 incomplays obtally perate as TH fellowith prosselp largers in transmission and are backlift.

the term Mist displays here cevers any matrix display to stabiling time-grated in mulinear elements, figure position the applies matrix, also displays containing parsite elements. Such as varieties or displays containing parsite elements.

Military layer in this type are particularly stituture in IT applications of we example problet TUS- or for highly incremation displayer in an automorphic or alternation metric to the displayer and alternations the automatic separate of the object of the displayer and the dasplayer free to the displayer are displayed from the automatic displayer free to the applicate accordingly of the object of the displayer free to the application of the displayer free to the application of the displayer from the displayer free to the application of the displayer from the displayer free to the displayer from the displayer free to the displayer free to the displayer from the displayer free to the displayer free to the displayer from the displayer free to the displayer free to

regardal numbers sense apply to provide the line of an Milthighly wing to interact in with the interior curiated in the display of him ountial preductance of very inpursant of a displayer will the factor have acted to be regardance walked over a ling gas withouten.

The disableantage of the MLD-TH limplays dust well hithert our iters to the accomparatively liw to transport latitudy on the viewing angle dependence on to the difficulty of senerating deep shades in these displays.

likere thus a stinger to be a great lemand to MLZ displays which have very high resistivity at the same time as a broad sperating temperature range, short response times and a low timeshold wolltade which can be one to produce warrout may chades.

It is an item of the invention to provide MLC displays land, on the ECR effect which do not have the abovement ned displayable or only it at the library with exercise and at the character time have many to the provide the country that

Fig. 1. Sharp in an assent in an district of instruction exists of all assentant actions the authorized property of a propert

Fig. 1. And the top of the control of experience of the control of the control

The mimins arreding to the invention has very terms while values in the reparetime thresholds relatively high values is the all ling but such at the same time very high values of the same time very high values is the same time very high values of the same time very high ver

Stre preferred emboirments are mentioned below:

a A medium whichir allord hally ourpoided one is more compounds if the formula II:

$$R^{2} \longrightarrow H \longrightarrow O \longrightarrow (O)CH_{2...}, \qquad II$$

 $_{
m P}$ is independently as defined for $_{
m P}$, P and R .

grand to the state which

y is from I to 3.


A greating with the additionally a superiore one or some terms of the contract of the contract

$$R^{2}$$
 A \rightarrow H \bullet R^{32}

1.11

Acres -

and an experimental and an experimental control of the second of the sec



- A mean model to reprove two three forms of the prejensibly two, three or into companion in the formula II.
 - A paint and the represent of least two graphs and in the formula is:
 - e Area do in which the propertion is try dolor to the formula II in the total mixture is at least I sy we dit preferably at least L by weight.
- 10 f A medium in which the proportion of region is of the formula in the total mixture in at least to by weight presentably at least 1. By weight.
- A medium in which the properties of compounds of the results in the total mixture in at least a ly we wint.
 - in A medium in which the proportion of someounds of the terminality of the terminality of the terminal decay.
 - 2 A feeting winter receive that Is an income typically as a modern ratio for all and a sub-like.

Particular prefetance in sixen to the tip under the tip and the tip and the contract of the co

alamiji di:

oslayi — .e a straight-shain Alkyl radigal having 1 - radi h atoma.

A medium who to admits makey why comes a some unimelected from the formulae live to IIId:

also entre esta

al Ryll and

aukylt – aras eart. Inaspendently to ne an ther a stragism main alkyl sadictel Cambro i to tables ar mai

The median are ring to the invention preferally specification of an invention of the formula transfer as a second of the contraction.

el ay neogir li ng a rae tag dair i tae taruladi

:1. 1

n " by weldh i ne fonde try unde i the rammia ii.

I A medium which addits nally comprised newspire on my unit in the institute.

$$F F$$

$$R H O O O H C_x H_{2x}.$$

$$F F F$$

$$R^3 - H O O O H C_x H_{2x}.$$

8 * 8,448 . **

Frank France reading independent by the new and increase and because the Frank Frank

The artists of the second of t

. He is a man which is a f⊕or . Neiligh or my a least that the second of the second o

 $R^{20} - H \leftarrow O - OCF_{2} - O - C_{1-aikyl}$ F = F $R^{21} - H \leftarrow H \leftarrow CF_{2}O - O - C_{1-aikyl}$ F = F $R^{22} - H \leftarrow H \leftarrow CCF_{2} - O - C_{1-aikyl}$

in which F E are each, independently is the another, as defined for R . F and R , and r and r are each, independently of one another, i.e., R is H, α H, α H or n-C-H.

i A medium in which the compound of the formula II is selected from the group consisting of flatter Ilg:

in which Process as defined as ve. and site 1 look is preparably straight chain alkylonaving of the votation atoms, which is alkenylog 3E-alkenylog.

- A medium which comprises one or more compounds of the remula lla and or lla.
- properties of the formula

la sent sa se la ciercia disenzia disenzia se disenzia se Sententantanta di la ciari sententa di la ciari disen

and contraints to grantly and attended to an about 100 and 100 from the district of the contract of the contra

The liquid strystal pixture preferreby has a separtic pixture range of at least (v_i) for an example of v_i

The relational when with γ is preferrebly γ where. In particular γ led means.

The stretch cases Δn in the lapted tryptal mixture is generally between 1.14 and 1.15, preferedary between 1.76 and 7.11, and we the dielectric constant ϵ - ϵi greater than 4: equal to 3. preferably from 3.5 is $\frac{1}{2}$.9.

The dielectrics may also coprise further additives which are known to the person skilled in the art and are described in the literature.

For example of the of ple-cinera dyes can be added, incrined the toplettime of a series preferably critical energy included by a series of the example of here expless salts of the whole of the example. Haller et al., Mol. Cryst. Liq. tryst., There example, pages 148-188 slows of are improving the confidentially of the confidential tryst. The confidential trial trial and the confidential trial trial

The Children like the present of the distribution of the little and the little an

nazed in grandard rethods which are bestrined in the

ine news of light type all numbers in the limplays according to the invention senseably of manage two transports with themselves trusted of one or more individual compounds.

rempinent A has significantly negative dislectric anisotropy and gives the negative phase a dislectric anisotropy \$i \$ 1.4. It preferably comprises compounds in the ferraless II and II.

The proportion of component A is preferably between 48 and 1:10, in particular between 31 and 1110.

For component A, one - r more individual compoundes having a $\Delta \epsilon \le -1.8$ are preferably selected. The smaller the proportion of component A in the total mixture, the more negative this value must be.

the problem is has problemed nematogeneity and a clew viscosity of not more than a one so, preferably not more than if mm s , at 1.0. It preferably compaises groupounds of the formulae ID and III.

Factional analysis presented that the action of a present of the properties of the p

demonstrate in the manufactor of enable topic to

• -

Introduction of the materials are known to the periodic relibed in the art from the liberature. Partyother preference in from to the construction in the formula 111.

 In audituum thege liquid trystal phases san aus symash muse than le sympthema, preferably from le to he companema.

The phases preferably someon from 4 to 10, in particular for 11, supposition of the compilae 11, 12, 11 and optionally 111.

Besides compounds of the formulae if, it, if and lift is also possible for other constituents to be present, for example in an arount of up to 45 of the total nixture, but preferably up to 35, in particular up to 3.

The other constituents are preferably selected from negatic or negaticente substances. In particular known adamances from the classes consisting of the anaxy persense, fembylidentamilines, applienyls, terphenyls, pienyl a cyclohexyl bencares, phenylcyclonexanes, cyclohexyl biphenyls, cyclohexylcyclonexanes, cyclohexylchinexylcyclonexanes, cyclohexylchinexylcyclonexanes, cyclohexylchinexylcyclonexanes, cyclohexylchinexylcyclonexanes, cyclohexylchinexylcyclonexanes, cyclohexylchinexy

ing stant tog omin who we have because an object which is Experi regularly mixtures on the type was like who were well as the or or its 100

grande gant anno 1964, in the state of the s

8 4

distance is a red of pricing in factory to make the type of members and type of the members of the state of t

1 28		
	11 24 -	*::::::::::::::::::::::::::::::::::::::
	•	111 111 (
	• 1,	31 <u>.2</u> _ 0 · .
	40 × 111 1, 4	

or a 3-0 single bond. It is halogen, preferably chlorine, or all and P and P are each alkyl, alkenyl, alkowy, alkanoyloxy or alkowycarbonyloxy having up to 18. preferably up to 8, parbon atoms, or one of these radicals is alternatively CM, NO, NO, NOS, CF, F, Cl or Bf.

In past of these compounds. P and P are different from me and these not these radicals usually seind an alkyloricalk-my stop. However, then variable of the proposed substituents are also conset. Many puch substances or mixtures thereof are commercially and labe. All these substances can be prepared by perhops which are an ordered.

In will be appreciated by a person skilled in the satthat the FTE mixture satisfains to the invention may also exprise companie in which is a example. He II, and I of Fibrurian replaced by the corresponding isotopes.

The contract of the contract o

Mith at further eld rate in it is delined that he satisfied in the art can unling the preferred in dearlight in whilther the present invention to the full wind preferred agention end there is an exhibit as reserved in the remainder of the discussions and that instanted of the remainder of the discussions in any way whatsuever.

in the foresting and in the islighted examples, all temperatures are set forth uncorrected in degrees deligion; and, unless otherwise individual, all parts and percentages are by weight.

The entire displashing of all applications, patents and publications of the appreciation of the application application DE 11. If the char, filed April 14, 1717, is hereby incorparated by referency.

Besides the compounds of the formulae II and II, the Intention II and III, the provide the invention preferably comprise one or more of the compounds rent. Selled w.

The relieving distribute are used: if (n, m = 1-8; g = 1 6)

F F F --- F

COPHOMES CHART H - H - O COLHAR.

CCP-nmFF C.H_{2...} H ← H ← O ← C_mH_{2m}.

CBC-nmf C.H. H → O - O — H → C.H.....

080 mm 0.-... 4 - 0 - 0 - H - 0.H...

oppropriate the Second of the

F F F F C.H.... H ← O ← O ← H ← C.H_{c...}. 0YY0-n-m FFFF CCYY-n- 0 m C,H_{2m} H ← H ← 0 ← C ← O)-C_H_{2m}. E E C.Y-n-m 00H-nm 0.H_{22.}... H ► ** H ► 0.,H_{24.}.. 3.-g., · d •-- - • 30 n . t F F QP.5 J.m.F.1 4 ج

$$CY-n-O1V$$
 C_nH_{2n+} H O $OCH_2CH=CH_2$

$$CY-n-OC(CH_3)=CH_2$$

$$CY-n-OC(CH_3)=CH_2$$

$$CY-n-OC(CH_3)=CH_3$$

$$CY-n-OC(CH_3)=CH_3$$

BCN-nm
$$C_nH_{2n+1}$$
 H H H C_mH_{2m+1}

COPC-rm
$$C_1H_{2-1}$$
 H C_mH_{2m} .

The abbreviations furthermore have the fellowing nearther:

er in the contract of the cont

 $\Delta n = -1$ figal anisatropy measured at 1 $^\circ$ C and 150 $^\circ$ mm

 Δc . The state of the state

egi. Eleginga set

y a manifest was noty resembled at 1 of phase)

eger johnsterner bit st

The company of the first section of the contract of the contra

Minigree energy, Seat

REAL COMMO

	- ·· ·	. · · 11	· · · · · · · · · · · · · · · · · · ·
EMERIARE	=	teating into the	
: 11	·· .	$\sum_{i=1}^{n} \frac{1}{n} \left(\frac{1}{n} - \frac{1}{n} \right) = \frac{1}{n} \left(\frac{1}{n} - \frac{1}{n} - \frac{1}{n} \right) = \frac{1}{n} \left(\frac{1}{n} - \frac{1}{n} - \frac{1}{n} \right) = \frac{1}{n} \left(\frac{1}{n} - \frac{1}{n} - \frac{1}{n} - \frac{1}{n} \right) = \frac{1}{n} \left(\frac{1}{n} - 1$	
• • • • • • • • • • • • • • • • • • •		∆ε in Riida a in i	:
	÷ . *	e *. kHn 1]:	• . •
	÷*	y indicated as the	*
, [31] 3 m 1		ild if rells: _ nem	
		A -1 C, 37 C, 41 C	

Examples ...

		₹) + II:	· : :
HUH - 4MM		dieding print i 🕌	* • •
		Δn [$n = \frac{1}{2} n = \frac{1}$	
BIH j	* •	Δv 11 kH2. 1 ω_{ij} :	
1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1		€ XHH ':	
1 1 1 1		lis in rells: nem. >	
11 Y (* 1.		At a fadi to	

Haranja ke d

1 11 × 4+11		ua · Mir	20)
1 11 + . FF	÷ .	Turkser Liber of Liber 1997	
1 10 4111		$\Delta m = m + m + m + m + m + m + m + m + m +$	
* ** . ***	8 .	Vicinity of the second of the second	
		4 4 4 4 4 4 4	٠.
		· · · · · · · · · · · · · · · · · · ·	:
e ere e		·*·	

W

			٠					
-				. *				7

[8 'H] + 4 [H]			•
		Trestar illia ja liita ja tiita	
e me i same	. " .	$\Delta : \mathbb{R}^{n-1} \to \mathbb{R}^{n-1} : \mathbb{R}^{n-1} \to \mathbb{R}^{n-1}$	
: 1 · EE	8 · ·	$\Delta \epsilon$, i.e., ϵ	E a
ZM KIPP		r (1 zHr, :	•
B B B B B B B B B B B B B B B B B B B		y ricercon to	*
* **			* •
· 11			
1111111	*		
1. 1			

Ем <u>ата</u> 1--

Politer	1 (g - N:	< ± ;
erte i læte	* .	Teaching point of the	
19 1 1 T = 1.1	* * *	Δn [1449 nm, 1.3 m]:	•
PCH i	(1)	$\Delta \epsilon$ [1 kHz, 1.1 k],:	
27 11 31	<u> </u>	ε ji kma n ma	
· · · · · · · · · · · · · · · · · · ·		y tuga en la	: *
·14		** :	
·; ·; ·			
:11 · LFF			

Emarty Lead

		Mean Mary miles	
ru · i		.\:\:\:\:\:\:\:\:\:\:\:\:\:\:\:\:\:\:\:	
		Δc , 1 with c . c :	·
	٠.	$\epsilon_{\rm cons}$. With	• • •
* ***			
	٠.	***	7.

hMari is

1 111 - 41 -		₹ · :::	. :
1 11 1-			
1 11 - 4 14		Δ :. \mathbb{R}^n . \mathbb{R}^n .	*
		$\Delta r = 1$ kills . The	·: ·
· · · · · · · · · · · · · · · · · · ·		e i kiin oo i, i	·
<u> </u>	* .	y miass. I is	
			: <u>.</u>
44			
NYP · . FF	* * * *		
	٠,		
Example:			
Pall alaks	± *\$	F + M:	
Man and Min		Treating limits of the	
I MI 4 JEE		Δi . \mathbb{R}^{n} is the same i . i .	• • • • • • • • • • • • • • • • • • • •
		$\Delta \epsilon$ leading in the	: .:
	100	€ .1 RHH. 1 - M1:	: 2
# '!!		y remains the	
	1 • .	***	
	*		
1 17			
Participals			
1311 - 5. 4Fb			
1 11 . P.			* .
		\ :.	
• • • • • • •		ΔC	
	• • •	e it eller i til	
		*** **	

Regular Ing I

F 18 + 488	- "	F + M:	
lotti olikk		Treating partitions that	
Fair Cale	. *	Δ :. The state of Δ :	
1,11 · 1,14 · 1		$\Delta \epsilon$, . which ; :	*
Hall t.	.	€ 1 kH+ 1]:	· · · · · ·
	-	Y market in the	
i /ii ·	٠.	is to	·
Fall GI o F			
·#11 1. 1.1,			

Hixtople 11

FOH : 4FF		\$) · N:	
iwii i imm	100	leating point of	<i>:</i> •
Vill 178		Δm , i ϵ ϵ m , i ϵ	
		$\Delta e^{-rac{\pi}{4}}$ 1 Wilm	. ·
		g l MHP, l t, :	• . •
BOH F.			
			. 15
17 .			
¥1 1			

With an Love . . .

: 1H + 1MF		Z + Mi	•
i tij sijy:		the graffich product	•
* ** 1 · · · · ·		\:. : :::: ::::::::::::::::::::::::::::	•
		As I will a light in the	• . •
		r laght in Ma	٠.
	• .	* :	
			* * *

Example 18

1 /H 1 FT	. ~ :	∴ : 11:	· .
: YE . 188		Territory of the	·
		Σ:. : : : :	
12-11-		$\Delta \epsilon$. The similar constants	· · · · · ·
	· .	€ 71 %HH. L 1 +	- , "
• 111 2 -	: .	0	· · · · · · · · · · · · · · · · · · ·
* * * * * * * * * * * * * * * * * * * *	• 0	y make the contract of the	- T
* 755	• • •	nir in wils: nem. +	
	3: 	4	
	8 .		

Emangile 14

POHRS. 4FF		S	
ión s afr		Heading into eq.:	
BC11 3		$\Delta n = \{ \cdot \mid \cdot \mid \cdot \mid n \in \mathbb{N} : $	*
		$\Delta \epsilon$ [1 AHL, ϵ [2]:	
		€ [] kHm, L. (2]:	
	• (
	•	y carlagest of the	
1 T .	· ·	nik n wiles net. v	

1 11 · · · · · · · · · · · · · · · · ·		₹ · iÎ:	•
1 11 1 1 1 1		Theoring in the	*
· 1		$\Delta : \mathbb{R}_{+} \times \mathbb{R}_{+} \times \mathbb{R}_{+}$ is	• 10 10 •
44		VC i dint, i :	•
ATT & J.		ε ji khiy ii ji;	
• • • • • • • • • • • • • • • • • • • •		:	
• • 1:	•	Table 1 to the control of the control of	
		article in the upon the second	

River 1 to 1

POH + SFF		thearthag and the tal	
rom der Strand		Δ:. · · · · · · : · · · · · · · · · · · ·	
PATE - APP			
		E LI RHA & Mi	
orgin s Ta		y nika (S). L. C.	
	· · · · ·	:	
HOH 31			
	* *-		
Example II			
POH-114PF		diwaring paunt (*1):	0 • 7
1 'H 4 LFF	·	And the first of the second of	
POH SEAPY		Δc [] sim, where	
OPT - L		ϵ [1 RHH, 1, 2]:	
JPQY-3 (IL	:	v (vii)	2.2
1 The state of the		y .mPa·s. Di Cl:	
· HYP Tale .	** .		
	" .		
	. · ·		
OCH 22	2		
Fix any less of			
ioni - 4111	- ·	Teahing print 1 ":	
7 H FF		Δz . The second z	
1 11 41 1	** * * * *	$\Delta v = 1$ William . The v	
0		€ I KMIL . Tit	
· · · · · · · · · · · · · · · · · · ·	•	÷ + + + + + + + + + + + + + + + + + + +	. :
÷ 11.	* .		
	*		

Harry et le

		tepartici poditi 1 1.4	
1 21 1 2 FF		<u> </u>	•
Family 4FF		$\Delta \epsilon$ (1 AHD D1 of):	· . · .
jejuje:	z	ϵ [1 kHd, ω] 204	÷ . 0
	- *		
й Ж		y rika ka a ba	
	•		
4	~ .		
elil te	: .		

Emaninas 4-

		clearing point 1 dis	- 5
PoH-1.4FF			
PCH #512FF	2.2.	Δn [E30 nm, 24 C]:	
PCH-514FF	40	$\Delta \epsilon$ [1] RH.1, III $^{\circ}$ [2]:	- 3 . 7
CCQY-3:11	- m	arepsilon [1] kHz, 2% Cl:	3.5
	5 4		1.14
124		y imparation of t	
• • • • • • • • • • • • • • • • • • • •			
	2		
2011			